



---

**ON A NEW CLASS OF EULERIAN'S TYPE INTEGRALS INVOLVING  
GENERALIZED HYPERGEOMETRIC FUNCTIONS**

SUNGTAE JUN, INSUK KIM\* AND ARJUN K. RATHIE

*Received 13 July, 2018; accepted 20 January, 2019; published 5 March, 2019.*

GENERAL EDUCATION INSTITUTE, KONKUK UNIVERSITY, CHUNGJU 380-701, REPUBLIC OF KOREA.

DEPARTMENT OF MATHEMATICS EDUCATION, WONKWANG UNIVERSITY, IKSAN, 570-749, REPUBLIC OF KOREA.

DEPARTMENT OF MATHEMATICS, VEDANT COLLEGE OF ENGINEERING AND TECHNOLOGY (RAJASTHAN TECHNICAL UNIVERSITY), BUNDI-323021, RAJASTHAN, INDIA.  
sjun@kku.ac.kr, iki@wku.ac.kr, arjunkumarrathie@gmail.com

**ABSTRACT.** Very recently Masjed-Jamei and Koepf established interesting and useful generalizations of various classical summation theorems for the  ${}_2F_1$ ,  ${}_3F_2$ ,  ${}_4F_3$ ,  ${}_5F_4$  and  ${}_6F_5$  generalized hypergeometric series. The main aim of this paper is to establish eleven Eulerian's type integrals involving generalized hypergeometric functions by employing these theorems. Several special cases have also been given.

*Key words and phrases:* Generalized hypergeometric functions; Classical summation theorems generalization.

*2000 Mathematics Subject Classification.* Primary 33C20, Secondary 33C05, 65B10.

## 1. INTRODUCTION

We define the well known Pochhammer symbol [7] for any complex number  $a$  as

$$(1.1) \quad (a)_n = \frac{\Gamma(a+n)}{\Gamma(a)} = \begin{cases} 1 & (n=0, a \in \mathbb{C} \setminus \{0\}) \\ a(a+1) \cdots (a+n-1) & (n \in \mathbb{N}, a \in \mathbb{C}), \end{cases}$$

where  $\Gamma(z)$  is the well known gamma function defined by

$$\Gamma(z) = \int_0^\infty e^{-x} x^{z-1} dx$$

for  $\operatorname{Re}(z) > 0$ .

Based on Pochhammer symbol (1.1), the generalized hypergeometric function with  $p$  numerator and  $q$  denominator parameters is defined [1, 3, 15, 18, 19] as

$$(1.2) \quad {}_pF_q \left[ \begin{matrix} a_1, & \dots, & a_p \\ b_1, & \dots, & b_q \end{matrix}; z \right] = \sum_{n=0}^{\infty} \frac{\prod_{i=1}^p (a_i)_n}{\prod_{i=1}^q (b_i)_n} \frac{z^n}{n!}$$

It can be easily verified by the ratio test [1, 2] that the series (1.2) is convergent for all  $p \leq q$ . Also it converges in  $|z| < 1$  for  $p = q+1$  and converges everywhere for  $p < q+1$  and converges nowhere ( $z \neq 0$ ) for  $p > q+1$ .

Further, if  $p = q+1$ , it converges absolutely for  $|z| = 1$  provided

$$\delta = \operatorname{Re} \left( \sum_{j=1}^q b_j - \sum_{j=1}^p b_j \right) > 0$$

holds and is conditionally convergent for  $|z| = 1$  and  $z \neq 1$  if  $-1 < \delta \leq 0$  and diverges for  $|z| = 1$  and  $z \neq 1$  if  $\delta \leq -1$ . For more details, we refer [15].

The two very important cases of the series (1.2), viz.

$${}_2F_1 \left[ \begin{matrix} a, & b \\ c \end{matrix}; z \right] = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!}$$

converges for  $|z| \leq 1$  and

$${}_1F_1 \left[ \begin{matrix} a \\ c \end{matrix}; z \right] = \sum_{n=0}^{\infty} \frac{(a)_n}{(c)_n} \frac{z^n}{n!}$$

converges everywhere, popularly known in the literature on the Gauss's hypergeometric function and Kummer's function or confluent hypergeometric function arising in many physical problems [12, 13, 17].

It is evident that whenever a generalized hypergeometric functions reduce to gamma function, the result are very important from the application point of view.

Moreover, here we shall mention the following classical summation theorems recorded in [1].

- Gauss Theorem for  $\operatorname{Re}(c - a - b) > 0$

$$(1.3) \quad {}_2F_1 \left[ \begin{matrix} a, & b \\ c \end{matrix}; 1 \right] = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)}$$

- Kummer's Theorem

$$(1.4) \quad {}_2F_1 \left[ \begin{matrix} a, & b \\ 1 + a - b \end{matrix} ; -1 \right] = \frac{\Gamma(1 + a - b)\Gamma(1 + \frac{1}{2}a)}{\Gamma(1 - b + \frac{1}{2}a)\Gamma(1 + a)}$$

- Second Gauss Theorem

$$(1.5) \quad {}_2F_1 \left[ \begin{matrix} a, & b \\ \frac{1}{2}(a + b + 1) \end{matrix} ; \frac{1}{2} \right] = \frac{\sqrt{\pi} \Gamma(\frac{1}{2}(a + b + 1))}{\Gamma(\frac{1}{2}(a + 1))\Gamma(\frac{1}{2}(b + 1))}$$

- Bailey's Theorem

$$(1.6) \quad {}_2F_1 \left[ \begin{matrix} a, & 1 - a \\ b \end{matrix} ; \frac{1}{2} \right] = \frac{\Gamma(\frac{1}{2}b)\Gamma(\frac{1}{2}(b + 1))}{\Gamma(\frac{1}{2}(a + b))\Gamma(\frac{1}{2}(b - a + 1))}$$

- Dixon's Theorem

$$(1.7) \quad {}_3F_2 \left[ \begin{matrix} a, & b, & c \\ 1 + a - b, & 1 + a - c \end{matrix} ; 1 \right] \\ = \frac{\Gamma(1 + \frac{1}{2}a)\Gamma(1 + a - b)\Gamma(1 + a - c)\Gamma(1 - b - c + \frac{1}{2}a)}{\Gamma(1 + a)\Gamma(1 - b + \frac{1}{2}a)\Gamma(1 - c + \frac{1}{2}a)\Gamma(1 + a - b - c)}$$

- Watson's Theorem

$$(1.8) \quad {}_3F_2 \left[ \begin{matrix} a, & b, & c \\ \frac{1}{2}(a + b + 1), & 2c \end{matrix} ; 1 \right] \\ = \frac{\sqrt{\pi} \Gamma(c + \frac{1}{2})\Gamma(\frac{1}{2}(a + b + 1))\Gamma(c - \frac{1}{2}(a + b - 1))}{\Gamma(\frac{1}{2}(a + 1))\Gamma(\frac{1}{2}(b + 1))\Gamma(c - \frac{1}{2}(a - 1))\Gamma(c - \frac{1}{2}(b - 1))}$$

- Whipple's Theorem

$$(1.9) \quad {}_3F_2 \left[ \begin{matrix} a, & 1 - a, & b \\ c, & 2b - c + 1 \end{matrix} ; 1 \right] \\ = \frac{\pi 2^{1-2b} \Gamma(c)\Gamma(2b - c + 1)}{\Gamma(\frac{1}{2}(a + c))\Gamma(b + \frac{1}{2}(a - c + 1))\Gamma(\frac{1}{2}(1 - a + c))\Gamma(b + 1 - \frac{1}{2}(a + c))}$$

- Pfaff-Saalschütz Theorem

$$(1.10) \quad {}_3F_2 \left[ \begin{matrix} a, & b, & -n \\ c, & 1 + a + b - c - n \end{matrix} ; 1 \right] = \frac{(c - a)_n(c - b)_n}{(c)_n(c - a - b)_n}$$

- Second Whipple's Theorem

$$(1.11) \quad {}_4F_3 \left[ \begin{matrix} a, & 1 + \frac{1}{2}a, & b, & c \\ \frac{1}{2}a, & a - b + 1, & a - c + 1 \end{matrix} ; -1 \right] = \frac{\Gamma(a - b + 1)\Gamma(a - c + 1)}{\Gamma(a + 1)\Gamma(a - b - c + 1)}$$

- Dougall's Theorem

$$(1.12) \quad {}_5F_4 \left[ \begin{matrix} a, & 1 + \frac{1}{2}a, & c, & d, & e \\ \frac{1}{2}a, & a - c + 1, & a - d + 1, & a - e + 1 \end{matrix} ; 1 \right] \\ = \frac{\Gamma(a - c + 1)\Gamma(a - d + 1)\Gamma(a - e + 1)\Gamma(a - c - d - e + 1)}{\Gamma(a + 1)\Gamma(a - d - e + 1)\Gamma(a - c - e + 1)\Gamma(a - c - d + 1)}$$

- Second Dougall's Theorem

$$(1.13) \quad {}_7F_6 \left[ \begin{matrix} a, & 1 + \frac{1}{2}a, & b, & c, & d, & 1 + 2a - b - c - d + n, & -n \\ \frac{1}{2}a, & a - b + 1, & a - c + 1, & a - d + 1, & b + c + d - a - n, & a + 1 + n \end{matrix} ; 1 \right] \\ = \frac{(a + 1)_n(a - b - c + 1)_n(a - b - d + 1)_n(a - c - d + 1)_n}{(a + 1 - b)_n(a + 1 - c)_n(a + 1 - d)_n(a + 1 - b - c - d)_n}$$

For finite sums of hypergeometric series, we will use the following symbol

$${}_pF_q^{(m)} \left[ \begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix}; z \right] = \sum_{n=0}^m \frac{\prod_{i=1}^p (a_i)_n}{\prod_{i=1}^q (b_i)_n} \frac{z^n}{n!},$$

where for instance

$${}_pF_q^{(-1)}(z) = 0, \quad {}_pF_q^{(0)}(z) = 1, \quad {}_pF_q^{(1)}(z) = 1 + \frac{a_1 \cdots a_p}{b_1 \cdots b_q} z.$$

By using the following relation [14].

$$\begin{aligned} (1.14) \quad & {}_pF_q \left[ \begin{matrix} a_1, \dots, a_{p-1}, 1 \\ b_1, \dots, b_{q-1}, m \end{matrix}; z \right] \\ &= \frac{\Gamma(b_1) \cdots \Gamma(b_{q-1}) \Gamma(a_1 - m + 1) \cdots \Gamma(a_{p-1} - m + 1)}{\Gamma(a_1) \cdots \Gamma(a_{p-1}) \Gamma(b_1 - m + 1) \cdots \Gamma(b_{q-1} - m + 1)} \frac{(m-1)!}{z^{m-1}} \\ &\quad \times \left\{ {}_{p-1}F_{q-1} \left[ \begin{matrix} a_1 - m + 1, \dots, a_{p-1} - m + 1 \\ b_1 - m + 1, \dots, b_{q-1} - m + 1 \end{matrix}; z \right] \right. \\ &\quad \left. - {}_{p-1}F_{q-1}^{(m-2)} \left[ \begin{matrix} a_1 - m + 1, \dots, a_{p-1} - m + 1 \\ b_1 - m + 1, \dots, b_{q-1} - m + 1 \end{matrix}; z \right] \right\}, \end{aligned}$$

very recently Masjed-Jamei and Koepf [11] have established generalizations of the classical summation theorems (1.3) to (1.13) in the following form:

$$\begin{aligned} (1.15) \quad & {}_3F_2 \left[ \begin{matrix} a, b, 1 \\ c, m \end{matrix}; 1 \right] \\ &= \frac{\Gamma(m)\Gamma(c)\Gamma(a-m+1)\Gamma(b-m+1)}{\Gamma(a)\Gamma(b)\Gamma(c-m+1)} \\ &\quad \times \left\{ \frac{\Gamma(c-m+1)\Gamma(c-a-b+m-1)}{\Gamma(c-a)\Gamma(c-b)} - {}_2F_1^{(m-2)} \left[ \begin{matrix} a-m+1, b-m+1 \\ c-m+1 \end{matrix}; 1 \right] \right\} \\ &= \Omega_1 \end{aligned}$$

$$\begin{aligned} (1.16) \quad & {}_3F_2 \left[ \begin{matrix} a, b, 1 \\ a-b+m, m \end{matrix}; -1 \right] \\ &= (-1)^{m-1} \frac{\Gamma(m)\Gamma(a-b+m)\Gamma(a-m+1)\Gamma(b-m+1)}{\Gamma(a)\Gamma(b)\Gamma(a-b+1)} \\ &\quad \times \left\{ \frac{\Gamma(a-b+1)\Gamma(1+\frac{1}{2}(a-m+1))}{\Gamma(2+a-m)\Gamma(m-b+\frac{1}{2}(a-m+1))} - {}_2F_1^{(m-2)} \left[ \begin{matrix} a-m+1, b-m+1 \\ a-b+1 \end{matrix}; -1 \right] \right\} \\ &= \Omega_2 \end{aligned}$$

$$\begin{aligned}
(1.17) \quad & {}_3F_2 \left[ \begin{matrix} a, b, 1 \\ \frac{1}{2}(a+b+1), m \end{matrix} ; \frac{1}{2} \right] \\
&= 2^{m-1} \frac{\Gamma(m)\Gamma(\frac{1}{2}(a+b+1))\Gamma(a-m+1)\Gamma(b-m+1)}{\Gamma(a)\Gamma(b)\Gamma(-m+1+\frac{1}{2}(a+b+1))} \\
&\times \left\{ \frac{\sqrt{\pi}\Gamma(-m+1+\frac{1}{2}(a+b+1))}{\Gamma(1+\frac{1}{2}(a-m))\Gamma(1+\frac{1}{2}(b-m))} - {}_2F_1^{(m-2)} \left[ \begin{matrix} a-m+1, b-m+1 \\ -m+1+\frac{1}{2}(a+b+1) \end{matrix} ; \frac{1}{2} \right] \right\} \\
&= \Omega_3
\end{aligned}$$

$$\begin{aligned}
(1.18) \quad & {}_3F_2 \left[ \begin{matrix} a, 2m-a-1, 1 \\ b, m \end{matrix} ; \frac{1}{2} \right] \\
&= 2^{m-1} \frac{\Gamma(m)\Gamma(b)\Gamma(a-m+1)\Gamma(m-a)}{\Gamma(a)\Gamma(2m-a-1)\Gamma(b-m+1)} \\
&\times \left\{ \frac{\Gamma(\frac{1}{2}(b-m+1))\Gamma(\frac{1}{2}(b-m+2))}{\Gamma(-m+1+\frac{1}{2}(a+b))\Gamma(\frac{1}{2}(b-a+1))} - {}_2F_1^{(m-2)} \left[ \begin{matrix} a-m+1, m-a \\ b-m+1 \end{matrix} ; \frac{1}{2} \right] \right\} \\
&= \Omega_4
\end{aligned}$$

$$\begin{aligned}
(1.19) \quad & {}_4F_3 \left[ \begin{matrix} a, b, c, 1 \\ a-b+m, a-c+m, m \end{matrix} ; 1 \right] \\
&= \frac{\Gamma(m)\Gamma(a-b+m)\Gamma(a-c+m)\Gamma(a+1-m)\Gamma(b+1-m)\Gamma(c+1-m)}{\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(a-b+1)\Gamma(a-c+1)} \\
&\times \left\{ \frac{\Gamma(\frac{1}{2}(a+3-m))\Gamma(a-b+1)\Gamma(a-c+1)\Gamma(-b-c+\frac{1}{2}(a+3m-1))}{\Gamma(a+2-m)\Gamma(-b+\frac{1}{2}(a+m+1))\Gamma(-c+\frac{1}{2}(a+m+1))\Gamma(a-b-c+m)} \right. \\
&\quad \left. - {}_3F_2^{(m-2)} \left[ \begin{matrix} a-m+1, b-m+1, c-m+1 \\ a-b+1, a-c+1 \end{matrix} ; 1 \right] \right\} \\
&= \Omega_5
\end{aligned}$$

$$\begin{aligned}
(1.20) \quad & {}_4F_3 \left[ \begin{matrix} a, b, c, 1 \\ \frac{1}{2}(a+b+1), 2c+1-m, m \end{matrix} ; 1 \right] \\
&= \frac{\Gamma(m)\Gamma(\frac{1}{2}(a+b+1))\Gamma(2c+1-m)\Gamma(a+1-m)\Gamma(b+1-m)\Gamma(c+1-m)}{\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(-m+\frac{1}{2}(a+b+3))\Gamma(2c-2m+2)} \\
&\times \left\{ \frac{\sqrt{\pi}\Gamma(c-m+\frac{3}{2})\Gamma(-m+\frac{1}{2}(a+b+3))\Gamma(c-\frac{1}{2}(a+b-1))}{\Gamma(1+\frac{1}{2}(a-m))\Gamma(1+\frac{1}{2}(b-m))\Gamma(c+1-\frac{1}{2}(a+m))\Gamma(c+1-\frac{1}{2}(b+m))} \right. \\
&\quad \left. - {}_3F_2^{(m-2)} \left[ \begin{matrix} a-m+1, b-m+1, c-m+1 \\ -m+1+\frac{1}{2}(a+b+1), 2c-2m+2 \end{matrix} ; 1 \right] \right\} \\
&= \Omega_6
\end{aligned}$$

$$\begin{aligned}
(1.21) \quad & {}_4F_3 \left[ \begin{matrix} a, & 2m-1-a, & b, & 1 \\ c, & 2b-c+1, & m & \end{matrix} ; 1 \right] \\
&= \frac{\Gamma(m)\Gamma(c)\Gamma(2b-c+1)\Gamma(m-a)\Gamma(a+1-m)\Gamma(b+1-m)}{\Gamma(a)\Gamma(b)\Gamma(2m-1-a)\Gamma(c+1-m)\Gamma(2b-c-m+2)} \\
&\quad \times \left\{ \frac{\pi 2^{2m-2b-1} \Gamma(c-m+1)}{\Gamma(-m+1+\frac{1}{2}(a+c))\Gamma(-m+1+b+\frac{1}{2}(a-c+1))\Gamma(\frac{1}{2}(1-a+c))} \right. \\
&\quad \times \left. \frac{\Gamma(2b-c-m+2)}{\Gamma(b+1-\frac{1}{2}(a+c))} - {}_3F_2 \left[ \begin{matrix} a-m+1, & b-m+1, & m-a \\ c-m+1, & 2b-c-m+2 & \end{matrix} ; 1 \right] \right\} \\
&= \Omega_7
\end{aligned}$$

$$\begin{aligned}
(1.22) \quad & {}_4F_3 \left[ \begin{matrix} a, & b, & -n+m-1, & 1 \\ c, & 1+a+b-c-n, & m & \end{matrix} ; 1 \right] = \frac{(m-1)!(1-c)_{m-1}}{(1-a)_{m-1}(1-b)_{m-1}} \\
&\quad \times \frac{(c-a-b+n)_{m-1}}{(n+2-m)_{m-1}} \times \left\{ \frac{(c-a)_n(c-b)_n}{(c+1-m)_n(c-a-b+m-1)_n} \right. \\
&\quad \left. - {}_3F_2 \left[ \begin{matrix} a-m+1, & b-m+1, & -n \\ c-m+1, & 2+a+b-c-m-n & \end{matrix} ; 1 \right] \right\} \\
&= \Omega_8
\end{aligned}$$

$$\begin{aligned}
(1.23) \quad & {}_5F_4 \left[ \begin{matrix} a, & \frac{1}{2}(a+m+1), & b, & c, & 1 \\ \frac{1}{2}(a+m-1), & a-b+m, & a-c+m, & m & \end{matrix} ; -1 \right] = (-1)^{m-1} \Gamma(m) \\
&\quad \times \frac{\Gamma(\frac{1}{2}(a+m-1))\Gamma(a-b+m)\Gamma(a-c+m)\Gamma(\frac{1}{2}(a-m+3))\Gamma(a-m+1)}{\Gamma(a)\Gamma(b)\Gamma(c)\Gamma(\frac{1}{2}(a+m+1))\Gamma(\frac{1}{2}(a-m+1))} \\
&\quad \times \frac{\Gamma(b+1-m)\Gamma(c+1-m)}{\Gamma(a-b+1)\Gamma(a-c+1)} \times \left\{ \frac{\Gamma(1+a-b)\Gamma(1+a-c)}{\Gamma(2-m+a)\Gamma(m+a-b-c)} \right. \\
&\quad \left. - {}_4F_3 \left[ \begin{matrix} a-m+1, & b-m+1, & \frac{1}{2}(a-m+3), & c-m+1 \\ \frac{1}{2}(a-m+1), & a-b+1, & a-c+1 & \end{matrix} ; -1 \right] \right\} \\
&= \Omega_9
\end{aligned}$$

$$\begin{aligned}
(1.24) \quad & {}_6F_5 \left[ \begin{matrix} a, & \frac{1}{2}(a+m+1), & c, & d, & e, & 1 \\ \frac{1}{2}(a+m-1), & a-c+m, & a-d+m, & a-e+m, & m & \end{matrix} ; 1 \right] \\
&= \frac{\Gamma(m)\Gamma(\frac{1}{2}(a+m-1))\Gamma(a-c+m)\Gamma(a-d+m)\Gamma(a-e+m)}{\Gamma(a-c+1)\Gamma(a-d+1)\Gamma(a-e+1)} \\
&\quad \times \frac{\Gamma(a-m+1)\Gamma(\frac{1}{2}(a-m+3))\Gamma(c+1-m)\Gamma(d+1-m)\Gamma(e+1-m)}{\Gamma(a)\Gamma(c)\Gamma(d)\Gamma(e)\Gamma(\frac{1}{2}(a+m+1))\Gamma(\frac{1}{2}(a-m+1))} \\
&\quad \times \left\{ \frac{\Gamma(a-c+1)\Gamma(a-d+1)\Gamma(a-e+1)\Gamma(a-c-d-e+2m-1)}{\Gamma(2-m+a)\Gamma(a-c-e+m)\Gamma(a-d-e+m)\Gamma(a-c-d+m)} \right. \\
&\quad \left. - {}_5F_4 \left[ \begin{matrix} a-m+1, & c-m+1, & \frac{1}{2}(a-m+3), & d-m+1, & e-m+1 \\ \frac{1}{2}(a-m+1), & a-c+1, & a-d+1, & a-e+1 & \end{matrix} ; 1 \right] \right\} \\
&= \Omega_{10}
\end{aligned}$$

(1.25)

$$\begin{aligned}
& {}_8F_7 \left[ \begin{matrix} a, \frac{1}{2}(a+m+1), b, c, d, 2a-b-c-d+2m-1+n, m-n-1, 1 \\ \frac{1}{2}(a+m-1), a-b+m, a-c+m, a-d+m, b+c+d-a+1-m-n, a+n+1, m \end{matrix} ; 1 \right] \\
&= (-1)^{m-1} (m-1)! \times \frac{(\frac{1}{2}(3-a-m))_{m-1} (1-a+b-m)_{m-1}}{(\frac{1}{2}(1-a-m))_{m-1} (1-a)_{m-1}} \\
&\times \frac{(1-a+c-m)_{m-1} (1-a+d-m)_{m-1} (m+n+a-b-c-d)_{m-1} (-a-n)_{m-1}}{(1-b)_{m-1} (1-c)_{m-1} (1-d)_{m-1} (b+c+d-2a+2-2m-n)_{m-1} (n+2-m)_{m-1}} \\
&\times \left\{ \frac{(a-m+2)_n (a-b-c+m)_n (a-b-d+m)_n (a-c-d+m)_n}{(a-b+1)_n (a-c+1)_n (a-d+1)_n (a-b-c-d+2m-1)_n} \right. \\
&- {}_7F_6 \left[ \begin{matrix} a-m+1, \frac{1}{2}(a-m+3), b-m+1, c-m+1, d-m+1, 2a-b-c-d+m+n, -n \\ \frac{1}{2}(a-m+1), a-b+1, a-c+1, a-d+1, b+c+d-a+2-2m-n, a-m+n+2 \end{matrix} ; 1 \right] \left. \right\} \\
&= \Omega_{11}
\end{aligned}$$

It is interesting to mention here that for  $m = 1$ , the results (1.15) to (1.25) reduce to the results (1.3) to (1.13) respectively. For other generalizations and extensions of the results (1.4) to (1.9), we refer [6, 8, 9, 10, 16].

The aim of this paper is to establish eleven Eulerian's type integrals involving generalized hypergeometric function by employing the summation theorems (1.15) to (1.25). Several new and interesting special cases have also been given.

## 2. EULERIAN'S TYPE INTEGRALS

The eleven new class of Eulerian's type integrals involving generalized hypergeometric functions to be established in the paper are given in the following theorems.

**Theorem 2.1.** For  $m \in \mathbb{N}$ ,  $Re(b) > 0$ ,  $Re(c-b) > 0$  and  $Re(c-a-b+m) > 1$ , the following result holds true.

$$(2.1) \quad \int_0^1 x^{b-1} (1-x)^{c-b-1} {}_2F_1 \left[ \begin{matrix} a, 1 \\ m \end{matrix} ; x \right] dx = \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} \Omega_1,$$

where  $\Omega_1$  is the same as given in (1.15).

*Proof.* For proving Theorem 2.1, denoting the left-hand side of (2.1) by  $I$ , expressing  ${}_2F_1$  as a series, change the order of integration and summation, evaluating the beta-integral, we have

$$I = \sum_{n=0}^{\infty} \frac{(a)_n (1)_n}{(m)_n n!} \frac{\Gamma(b+n)\Gamma(c-b)}{\Gamma(c+n)}.$$

Using (1.1) and after some simplification, we have

$$I = \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} \sum_{n=0}^{\infty} \frac{(a)_n (b)_n (1)_n}{(c)_n (m)_n} \frac{1}{n!}.$$

Summing up the series, we finally get

$$I = \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} {}_3F_2 \left[ \begin{matrix} a, b, 1 \\ c, m \end{matrix} ; 1 \right].$$

Now, we observe that the  ${}_3F_2$  appearing can be evaluated with the help of the result (1.15) and we easily arrive at the right side of (2.1). This completes the proof of Theorem 2.1.

■

**Corollary 2.2.** *In Theorem 2.1, if we take  $m = 2, 3$  (omitting the trivial case of  $m = 1$ ), we respectively get the following integrals.*

$$\begin{aligned} & \int_0^1 x^{b-1}(1-x)^{c-b-1} {}_2F_1 \left[ \begin{matrix} a, & 1 \\ & 2 \end{matrix}; x \right] dx \\ &= \frac{(c-1)}{(a-1)(b-1)} \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} \left\{ \frac{\Gamma(c-1)\Gamma(c-a-b+1)}{\Gamma(c-a)\Gamma(c-b)} - 1 \right\} \end{aligned}$$

and

$$\begin{aligned} & \int_0^1 x^{b-1}(1-x)^{c-b-1} {}_2F_1 \left[ \begin{matrix} a, & 1 \\ & 3 \end{matrix}; x \right] dx \\ &= \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} \frac{2(c-2)_2}{(a-2)_2(b-2)_2} \\ & \quad \times \left\{ \frac{\Gamma(c-2)\Gamma(c-a-b+2)}{\Gamma(c-a)\Gamma(c-b)} - \frac{ab+c-2a-2b+2}{c-2} \right\}. \end{aligned}$$

The following Theorems and the corresponding Corollaries 2.3 to 2.22 can be obtained by employing the results (1.16) to (1.25). Hence, they are given here without proof.

**Theorem 2.3.** *For  $m \in \mathbb{N}$ ,  $\operatorname{Re}(b) > 0$  and  $\operatorname{Re}(a - 2b + m) > 0$ , the following result holds true.*

$$(2.2) \quad \int_0^1 x^{b-1}(1-x)^{a-2b+m-1} {}_2F_1 \left[ \begin{matrix} a, & 1 \\ & m \end{matrix}; -x \right] dx = \frac{\Gamma(b)\Gamma(a-2b+m)}{\Gamma(a-b+m)} \Omega_2,$$

where  $\Omega_2$  is the same as given in (1.16).

**Corollary 2.4.** *In Theorem 2.3, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.*

$$\begin{aligned} & \int_0^1 x^{b-1}(1-x)^{a-2b}(1+x)^{-a} dx = \frac{\Gamma(b)\Gamma(1+a-2b)\Gamma(1+\frac{1}{2}a)}{\Gamma(1+a)\Gamma(1+\frac{1}{2}a-b)}, \\ & \int_0^1 x^{b-1}(1-x)^{a-2b+1} {}_2F_1 \left[ \begin{matrix} a, & 1 \\ & 2 \end{matrix}; x \right] dx \\ &= \frac{(a-b+1)}{(a-1)(b-1)} \frac{\Gamma(b)\Gamma(a-2b+2)}{\Gamma(a-b+2)} \left\{ 1 - \frac{\Gamma(1+a-b)\Gamma(\frac{1}{2}a+\frac{1}{2})}{\Gamma(a)\Gamma(\frac{1}{2}a-b+\frac{3}{2})} \right\} \end{aligned}$$

and

$$\begin{aligned} & \int_0^1 x^{b-1}(1-x)^{a-2b+2} {}_2F_1 \left[ \begin{matrix} a, & 1 \\ & 3 \end{matrix}; x \right] dx \\ &= \frac{2(a-b+1)_2}{(a-2)_2(b-2)_2} \frac{\Gamma(b)\Gamma(a-2b+3)}{\Gamma(a-b+3)} \\ & \quad \times \left\{ \frac{\Gamma(\frac{1}{2}a)\Gamma(1+a-b)}{\Gamma(a-1)\Gamma(\frac{1}{2}a-b+2)} - \frac{3a+b-ab-3}{1+a-b} \right\}. \end{aligned}$$

**Theorem 2.5.** *For  $m \in \mathbb{N}$ ,  $\operatorname{Re}(b) > 0$  and  $\operatorname{Re}(a - b + 1) > 0$ , the following result holds true.*

$$(2.3) \quad \int_0^1 x^{b-1}(1-x)^{\frac{1}{2}(a-b-1)} {}_2F_1 \left[ \begin{matrix} a, & 1 \\ & m \end{matrix}; \frac{1}{2}x \right] dx = \frac{\Gamma(b)\Gamma(\frac{1}{2}(a-b+1))}{\Gamma(\frac{1}{2}(a+b+1))} \Omega_3,$$

where  $\Omega_3$  is the same as given in (1.17).



**Corollary 2.6.** *In Theorem 2.5, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.*

$$\int_0^1 x^{b-1}(1-x)^{\frac{1}{2}(a-b-1)}\left(1-\frac{1}{2}x\right)^{-a}dx = \frac{\Gamma(\frac{1}{2})\Gamma(b)\Gamma(\frac{1}{2}(a-b+1))}{\Gamma(\frac{1}{2}(a+1))\Gamma(\frac{1}{2}(b+1))},$$

$$\int_0^1 x^{b-1}(1-x)^{\frac{1}{2}(a-b-1)}{}_2F_1\left[\begin{matrix} a, & 1 \\ 2 & \end{matrix}; \frac{1}{2}x\right]dx$$

$$= \frac{(a+b-1)}{(a-1)(b-1)} \frac{\Gamma(b)\Gamma(\frac{1}{2}(a-b+1))}{\Gamma(\frac{1}{2}(a+b+1))} \left\{ \frac{\sqrt{\pi}\Gamma(\frac{1}{2}(a+b-1))}{\Gamma(\frac{1}{2}a)\Gamma(\frac{1}{2}b)} - 1 \right\}$$

and

$$\int_0^1 x^{b-1}(1-x)^{\frac{1}{2}(a-b-1)}{}_2F_1\left[\begin{matrix} a, & 1 \\ 3 & \end{matrix}; \frac{1}{2}x\right]dx$$

$$= \frac{2(a+b-1)(a+b-3)}{(a-2)_2(b-2)_2} \frac{\Gamma(b)\Gamma(\frac{1}{2}(a-b+1))}{\Gamma(\frac{1}{2}(a+b+1))}$$

$$\times \left\{ \frac{\sqrt{\pi}\Gamma(\frac{1}{2}(a+b-3))}{\Gamma(\frac{1}{2}(a-1))\Gamma(\frac{1}{2}(b-1))} - \frac{ab-a-b+1}{a+b-3} \right\}.$$

**Theorem 2.7.** *For  $m \in \mathbb{N}$ ,  $Re(a) > 0$  and  $Re(b-a) > 0$ , the following result holds true.*

$$(2.4) \quad \int_0^1 x^{a-1}(1-x)^{b-a-1}{}_2F_1\left[\begin{matrix} 2m-a-1, & 1 \\ m & \end{matrix}; \frac{1}{2}x\right]dx = \frac{\Gamma(a)\Gamma(b-a)}{\Gamma(b)}\Omega_4,$$

where  $\Omega_4$  is the same as given in (1.18).

**Corollary 2.8.** *In Theorem 2.7, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.*

$$\int_0^1 x^{a-1}(1-x)^{b-a-1}\left(1-\frac{1}{2}x\right)^{-a-1}dx = \frac{2^{1-b}\Gamma(\frac{1}{2})\Gamma(a)\Gamma(b-a)}{\Gamma(\frac{1}{2}(a+b))\Gamma(\frac{1}{2}(b-a+1))},$$

$$\int_0^1 x^{a-1}(1-x)^{b-a-1}{}_2F_1\left[\begin{matrix} 3-a, & 1 \\ 2 & \end{matrix}; \frac{1}{2}x\right]dx$$

$$= \frac{2(1-b)}{(1-a)_2} \frac{\Gamma(a)\Gamma(b-a)}{\Gamma(b)} \left\{ \frac{\Gamma(\frac{1}{2}(b-1))\Gamma(\frac{1}{2}b)}{\Gamma(\frac{1}{2}(a+b)-1)\Gamma(\frac{1}{2}(b-a+1))} - 1 \right\}$$

and

$$\int_0^1 x^{a-1}(1-x)^{b-a-1}{}_2F_1\left[\begin{matrix} 5-a, & 1 \\ 3 & \end{matrix}; \frac{1}{2}x\right]dx$$

$$= \frac{8(b-2)_2}{(a-4)_4} \frac{\Gamma(a)\Gamma(b-a)}{\Gamma(b)} \left\{ \frac{\Gamma(\frac{1}{2}(b-1))\Gamma(\frac{1}{2}(b-2))}{\Gamma(\frac{1}{2}(a+b)-2)\Gamma(\frac{1}{2}(b-a+1))} - \frac{5a-a^2+2b-10}{2(b-2)} \right\}.$$

**Theorem 2.9.** *For  $m \in \mathbb{N}$ ,  $Re(c) > 0$  and  $Re(a-2c+m) > 0$ , the following result holds true.*

$$(2.5) \quad \int_0^1 x^{c-1}(1-x)^{a-2c+m-1}{}_3F_2\left[\begin{matrix} a, & b, & 1 \\ a-b+m, & m & \end{matrix}; x\right]dx = \frac{\Gamma(c)\Gamma(a-2c+m)}{\Gamma(a-c+m)}\Omega_5,$$

where  $\Omega_5$  is the same as given in (1.19).

**Corollary 2.10.** *In Theorem 2.9, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.*

$$\begin{aligned} & \int_0^1 x^{c-1}(1-x)^{a-2c} {}_2F_1 \left[ \begin{matrix} a, & b \\ 1+a-b \end{matrix}; x \right] dx \\ &= \frac{\Gamma(c)\Gamma(1+a-2c)\Gamma(1+\frac{1}{2}a)\Gamma(1+a-b)\Gamma(1+\frac{1}{2}a-b-c)}{\Gamma(1+a)\Gamma(1+\frac{1}{2}a-b)\Gamma(1+\frac{1}{2}a-c)\Gamma(1+a-b-c)}, \\ & \int_0^1 x^{c-1}(1-x)^{a-2c+1} {}_3F_2 \left[ \begin{matrix} a, & b, & 1 \\ a-b+2, & 2 \end{matrix}; x \right] dx \\ &= \frac{(1+a-b)(1+a-c)}{(a-1)(b-1)(c-1)} \frac{\Gamma(c)\Gamma(2+a-2c)}{\Gamma(a-c+2)} \\ & \quad \times \left\{ \frac{\Gamma(\frac{1}{2}(a+1))\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(\frac{1}{2}a-b-c+\frac{5}{2})}{\Gamma(a)\Gamma(\frac{1}{2}a-b+\frac{3}{2})\Gamma(\frac{1}{2}a-c+\frac{3}{2})\Gamma(2+a-b-c)} - 1 \right\} \end{aligned}$$

and

$$\begin{aligned} & \int_0^1 x^{c-1}(1-x)^{a-2c+2} {}_3F_2 \left[ \begin{matrix} a, & b, & c \\ a-b+3, & 3 \end{matrix}; x \right] dx \\ &= \frac{2(a-b+1)_2(a-c+1)_2}{(a-2)_2(b-2)_2(c-2)_2} \frac{\Gamma(c)\Gamma(a-2c+3)}{\Gamma(a-c+3)} \\ & \quad \times \left\{ \frac{\Gamma(\frac{1}{2}a)\Gamma(1+a-b)\Gamma(1+a-c)\Gamma(\frac{1}{2}a-b-c+4)}{\Gamma(a-1)\Gamma(\frac{1}{2}a-b+2)\Gamma(\frac{1}{2}a-c+2)\Gamma(3+a-b-c)} - \frac{(a-2)(b-2)(c-2)}{(a-b+1)(a-c+1)} - 1 \right\}. \end{aligned}$$

**Remark 2.1.** The first result given in the Corollary 2.10 is also recorded in [4, 5].

**Theorem 2.11.** *For  $m \in \mathbb{N}$  and  $\operatorname{Re}(c) > 0$ , the following result holds true.*

$$(2.6) \quad \int_0^1 x^{c-1}(1-x)^{c-m} {}_3F_2 \left[ \begin{matrix} a, & b, & 1 \\ \frac{1}{2}(a+b+1), & m \end{matrix}; x \right] dx = \frac{\Gamma(c)\Gamma(c-m+1)}{\Gamma(2c-m+1)} \Omega_6,$$

where  $\Omega_6$  is the same as given in (1.20).

**Corollary 2.12.** *In Theorem 2.11, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.*

$$\begin{aligned} & \int_0^1 x^{c-1}(1-x)^{c-1} {}_2F_1 \left[ \begin{matrix} a, & b \\ \frac{1}{2}(a+b+1) \end{matrix}; x \right] dx \\ &= \frac{\Gamma(\frac{1}{2})\Gamma(c)\Gamma(c)\Gamma(c+\frac{1}{2})\Gamma(\frac{1}{2}(a+b+1))\Gamma(c-\frac{1}{2}(a+b-1))}{\Gamma(2c)\Gamma(\frac{1}{2}(a+1))\Gamma(\frac{1}{2}(b+1))\Gamma(c-\frac{1}{2}(a-1))\Gamma(c-\frac{1}{2}(b-1))}, \\ & \int_0^1 x^{c-1}(1-x)^{c-2} {}_3F_2 \left[ \begin{matrix} a, & b, & 1 \\ \frac{1}{2}(a+b+1), & 2 \end{matrix}; x \right] dx \\ &= \frac{(a+b-1)}{(a-1)(b-1)} \frac{\Gamma(c)\Gamma(c-1)}{\Gamma(2c-1)} \left\{ \frac{\sqrt{\pi}\Gamma(c-\frac{1}{2})\Gamma(\frac{1}{2}(a+b-1))\Gamma(c-\frac{1}{2}(a+b-1))}{\Gamma(\frac{1}{2}a)\Gamma(\frac{1}{2}b)\Gamma(c-\frac{1}{2}a)\Gamma(c-\frac{1}{2}b)} - 1 \right\} \end{aligned}$$

and

$$\int_0^1 x^{c-1}(1-x)^{c-3} {}_3F_2 \left[ \begin{matrix} a, & b, & 1 \\ \frac{1}{2}(a+b+1), & 3 \end{matrix} ; x \right] dx$$

$$= \frac{(2c-3)(a+b-1)(a+b-3)}{(c-1)(a-2)_2(b-2)_2} \frac{\Gamma(c)\Gamma(c-2)}{\Gamma(2c-2)}$$

$$\times \left\{ \frac{\sqrt{\pi}\Gamma(c-\frac{3}{2})\Gamma(\frac{1}{2}(a+b-3))\Gamma(c-\frac{1}{2}(a+b-1))}{\Gamma(\frac{1}{2}(a-1))\Gamma(\frac{1}{2}(b-1))\Gamma(c-\frac{1}{2}(a+1))\Gamma(c-\frac{1}{2}(b+1))} - \frac{(a-2)(b-2)}{a+b-3} - 1 \right\}.$$

**Remark 2.2.** The first result given in the Corollary 2.12 is also recorded in [4, 5].

**Theorem 2.13.** For  $m \in \mathbb{N}$ ,  $Re(a) > 0$  and  $Re(c-a) > 0$ , the following result holds true.

$$(2.7) \quad \int_0^1 x^{a-1}(1-x)^{c-a-1} {}_3F_2 \left[ \begin{matrix} 2m-1-a, & b, & 1 \\ 2b-c+1, & m \end{matrix} ; x \right] dx = \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)} \Omega_7,$$

where  $\Omega_7$  is the same as given in (1.21).

**Corollary 2.14.** In Theorem 2.13, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.

$$\int_0^1 x^{a-1}(1-x)^{c-a-1} {}_2F_1 \left[ \begin{matrix} 1-a, & b \\ 2b-c+1 \end{matrix} ; x \right] dx$$

$$= \frac{\pi 2^{1-2b} \Gamma(a)\Gamma(c-a)\Gamma(2b-c+1)}{\Gamma(\frac{1}{2}(a+c))\Gamma(b+\frac{1}{2}(a-c+1))\Gamma(\frac{1}{2}(1-a+c))\Gamma(b+1-\frac{1}{2}(a+c))},$$

$$\int_0^1 x^{a-1}(1-x)^{c-a-1} {}_3F_2 \left[ \begin{matrix} 3-a, & b, & 1 \\ 2b-c+1, & 2 \end{matrix} ; x \right] dx$$

$$= \frac{(c-1)(c-2b)}{(a-2)_2(b-1)} \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)}$$

$$\times \left\{ \frac{\pi 2^{3-2b} \Gamma(c-1)\Gamma(2b-c)}{\Gamma(\frac{1}{2}(a+c)-1)\Gamma(b+\frac{1}{2}(a-c-1))\Gamma(\frac{1}{2}(1-a+c))\Gamma(b+1-\frac{1}{2}(a+c))} - 1 \right\}$$

and

$$\int_0^1 x^{a-1}(1-x)^{c-a-1} {}_3F_2 \left[ \begin{matrix} 5-a, & b, & 1 \\ 2b-c+1, & 3 \end{matrix} ; x \right] dx = \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)}$$

$$\times \left\{ \frac{\pi 2^{5-2b} \Gamma(c-2)\Gamma(2b-c+1)}{\Gamma(\frac{1}{2}(a+c)-2)\Gamma(b+\frac{1}{2}(a-c-3))\Gamma(\frac{1}{2}(1-a+c))\Gamma(b+1-\frac{1}{2}(a+c))} - \frac{(a-2)(3-a)(b-2)}{(c-2)(2b-c-1)} - 1 \right\}.$$

**Remark 2.3.** The first result given in the Corollary 2.14 is also recorded in [4, 5].

**Theorem 2.15.** For  $m \in \mathbb{N}$ ,  $Re(a) > 0$  and  $Re(c-a) > 0$ , the following result holds true.

$$(2.8) \quad \int_0^1 x^{a-1}(1-x)^{c-a-1} {}_3F_2 \left[ \begin{matrix} -n+m-1, & b, & 1 \\ 1+a+b-c-n, & m \end{matrix} ; x \right] dx = \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)} \Omega_8,$$

where  $\Omega_8$  is the same as given in (1.22).

**Corollary 2.16.** *In Theorem 2.15, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.*

$$\int_0^1 x^{a-1}(1-x)^{c-a-1} {}_2F_1 \left[ \begin{matrix} -n, & b \\ 1+a+b-c-n \end{matrix} ; x \right] dx = \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)} \frac{(c-a)_n(c-b)_n}{(c)_n(c-a-b)_n},$$

$$\int_0^1 x^{a-1}(1-x)^{c-a-1} {}_3F_2 \left[ \begin{matrix} -n+1, & b, & 1 \\ 1+a+b-c-n, & 2 \end{matrix} ; x \right] dx$$

$$= \frac{(1-c)(c-a-b+n)}{n(1-a)(1-b)} \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)} \times \left\{ \frac{(c-a)_n(c-b)_n}{(c)_n(c-a-b+1)_n} - 1 \right\}$$

and

$$\int_0^1 x^{a-1}(1-x)^{c-a-1} {}_3F_2 \left[ \begin{matrix} -n+2, & b, & 1 \\ 1+a+b-c-n, & 3 \end{matrix} ; x \right] dx$$

$$= \frac{2(1-c)_2(c-a-b+n)_2}{n(1-a)_2(1-b)_2} \frac{\Gamma(a)\Gamma(c-a)}{\Gamma(c)}$$

$$\times \left\{ \frac{(c-a)_n(c-b)_n}{(c-2)_n(c-a-b+2)_n} + \frac{n(a-2)(b-2)}{(c-2)(a+b-c-n-1)} - 1 \right\}.$$

**Theorem 2.17.** *For  $m \in \mathbb{N}$ ,  $\operatorname{Re}(b) > 0$  and  $\operatorname{Re}(a-2b+m) > 0$ , the following result holds true.*

$$(2.9) \quad \int_0^1 x^{b-1}(1-x)^{a-2b+m-1} {}_4F_3 \left[ \begin{matrix} a, & \frac{1}{2}(a+m+1), & c, & 1 \\ \frac{1}{2}(a+m-1), & a-c+m, & m \end{matrix} ; -x \right] dx$$

$$= \frac{\Gamma(b)\Gamma(a-2b+m)}{\Gamma(a-b+m)} \Omega_9,$$

where  $\Omega_9$  is the same as given in (1.23).

**Corollary 2.18.** *In Theorem 2.17, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.*

$$\int_0^1 x^{b-1}(1-x)^{a-2b} {}_3F_2 \left[ \begin{matrix} a, & \frac{1}{2}(a+2), & c \\ \frac{1}{2}a, & a-c+1 \end{matrix} ; -x \right] dx$$

$$= \frac{\Gamma(b)\Gamma(1+a-c)\Gamma(1+a-2b)}{\Gamma(1+a)\Gamma(1+a-b-c)},$$

$$\int_0^1 x^{b-1}(1-x)^{a-2b+1} {}_4F_3 \left[ \begin{matrix} a, & \frac{1}{2}(a+3), & c, & 1 \\ \frac{1}{2}(a+1), & a-c+2, & 2 \end{matrix} ; -x \right] dx$$

$$= \frac{(1+a-b)(1+a-c)}{(a+1)(b-1)(c-1)} \frac{\Gamma(b)\Gamma(a-2b+2)}{\Gamma(2+a-b)} \times \left\{ 1 - \frac{\Gamma(1+a-b)\Gamma(1+a-c)}{\Gamma(a)\Gamma(2+a-b-c)} \right\}$$

and

$$\int_0^1 x^{b-1}(1-x)^{a-2b+2} {}_4F_3 \left[ \begin{matrix} a, & \frac{1}{2}(a+4), & c, & 1 \\ \frac{1}{2}(a+3), & a-c+3, & 3 \end{matrix} ; -x \right] dx$$

$$= \frac{2(1+a-b)_2(1+a-c)_2}{(a+2)(a-1)(b-2)_2(c-2)_2} \frac{\Gamma(b)\Gamma(a-2b+3)}{\Gamma(3+a-b)}$$

$$\times \left\{ \frac{\Gamma(1+a-b)\Gamma(1+a-c)}{\Gamma(a-1)\Gamma(3+a-b-c)} + \frac{a(b-2)(c-2)}{(1+a-b)(1+a-c)} - 1 \right\}.$$

**Theorem 2.19.** For  $m \in \mathbb{N}$ ,  $\operatorname{Re}(b) > 0$  and  $\operatorname{Re}(a - 2c + m) > 0$ , the following result holds true.

$$(2.10) \quad \int_0^1 x^{c-1}(1-x)^{a-2c+m-1} {}_5F_4 \left[ \begin{matrix} a, \frac{1}{2}(a+m+1), d, e, 1 \\ \frac{1}{2}(a+m-1), a-d+m, a-e+m, m \end{matrix}; x \right] dx \\ = \frac{\Gamma(c)\Gamma(a-2c+m)}{\Gamma(a-c+m)} \Omega_{10},$$

where  $\Omega_{10}$  is the same as given in (1.24).

**Corollary 2.20.** In Theorem 2.19, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.

$$\int_0^1 x^{c-1}(1-x)^{a-2c} {}_4F_3 \left[ \begin{matrix} a, \frac{1}{2}(a+2), d, e \\ \frac{1}{2}a, a-d+1, a-e+1 \end{matrix}; x \right] dx \\ = \frac{\Gamma(c)\Gamma(a-2c)\Gamma(1+a-d)\Gamma(1+a-e)\Gamma(1+a-c-d-e)}{\Gamma(1+a)\Gamma(1+a-d-e)\Gamma(1+a-c-e)\Gamma(1+a-c-d)},$$

$$\int_0^1 x^{c-1}(1-x)^{a-2c+1} {}_5F_4 \left[ \begin{matrix} a, \frac{1}{2}(a+3), d, e, 1 \\ \frac{1}{2}(a+1), a-d+2, a-e+2, 2 \end{matrix}; x \right] dx \\ = \frac{(1+a-c)(1+a-d)(1+a-e)}{(1+a)(c-1)(d-1)(e-1)} \frac{\Gamma(c)\Gamma(a-2c+2)}{\Gamma(a-c+2)} \\ \times \left\{ \frac{\Gamma(1+a-c)\Gamma(1+a-d)\Gamma(1+a-e)\Gamma(3+a-c-d-e)}{\Gamma(a)\Gamma(2+a-d-e)\Gamma(2+a-c-e)\Gamma(2+a-c-d)} - 1 \right\}$$

and

$$\int_0^1 x^{c-1}(1-x)^{a-2c+2} {}_5F_4 \left[ \begin{matrix} a, \frac{1}{2}(a+4), d, e, 1 \\ \frac{1}{2}(a+2), a-d+3, a-e+3, 3 \end{matrix}; x \right] dx \\ = \frac{2(1+a-c)_2(1+a-d)_2(1+a-e)_2}{(a-1)(a+2)(c-2)_2(d-2)_2(e-2)_2} \frac{\Gamma(c)\Gamma(a-2c+3)}{\Gamma(a-c+3)} \\ \times \left\{ \frac{\Gamma(1+a-c)\Gamma(1+a-d)\Gamma(1+a-e)\Gamma(5+a-c-d-e)}{\Gamma(a-1)\Gamma(3+a-d-e)\Gamma(3+a-c-e)\Gamma(3+a-c-d)} \right. \\ \left. - \frac{a(c-2)(d-2)(e-2)}{(1+a-c)(1+a-d)(1+a-e)} \right\}.$$

**Theorem 2.21.** For  $m \in \mathbb{N}$ ,  $\operatorname{Re}(b) > 0$  and  $\operatorname{Re}(a - 2b + m) > 0$ , the following result holds true.

$$(2.11) \quad \int_0^1 x^{b-1}(1-x)^{a-2b+m-1} \\ \times {}_7F_6 \left[ \begin{matrix} a, \frac{1}{2}(a+m+1), c, d, 2a-b-c-d+2m+n-1, m-n-1 \\ \frac{1}{2}(a+m-1), a-c+m, a-d+m, b+c+d-a-m-n+1, a+n+1, m \end{matrix}; x \right] dx \\ = \frac{\Gamma(b)\Gamma(a-2b+m)}{\Gamma(a-b+m)} \Omega_{11},$$

where  $\Omega_{11}$  is the same as given in (1.25).

**Corollary 2.22.** In Theorem 2.21, if we take  $m = 1, 2, 3$ , we respectively get the following integrals.

$$\begin{aligned} & \int_0^1 x^{b-1}(1-x)^{a-2b} \\ & \times {}_6F_5 \left[ \begin{matrix} a, & \frac{1}{2}(a+2), & c, & d, & 2a-b-c-d+n+1, & -n \\ \frac{1}{2}a, & 1+a-c, & 1+a-d, & b+c+d-a-n, & a+n+1 \end{matrix} ; x \right] dx \\ & = \frac{\Gamma(b)\Gamma(a-2b+1)}{\Gamma(a-b+1)} \frac{(1+a)_n(a-b-c+1)_n(a-b-d+1)_n(a-c-d+1)_n}{(1+a-b)_n(1+a-c)_n(1+a-d)_n(1+a-b-c-d)_n}, \end{aligned}$$

$$\begin{aligned} & \int_0^1 x^{b-1}(1-x)^{a-2b+1} \\ & \times {}_7F_6 \left[ \begin{matrix} a, & \frac{1}{2}(a+3), & c, & d, & 2a-b-c-d+n+3, & 1-n, & 1 \\ \frac{1}{2}(a+1), & 1+a-c, & 1+a-d, & b+c+d-a-n-1, & a+n+1, & 2 \end{matrix} ; x \right] dx \\ & = \frac{(b-a-1)(c-a-1)(d-a-1)(n+2+a-b-c-d)(a+n)\Gamma(b)\Gamma(a-2b+1)}{n(1+a)(1-b)(1-c)(1-d)(b+c+d-2a-2-n)\Gamma(a-b+1)} \\ & \times \left\{ 1 - \frac{(a)_n(a-b-c+2)_n(a-b-d+2)_n(a-c-d+2)_n}{(1+a-b)_n(1+a-c)_n(1+a-d)_n(3+a-b-c-d)_n} \right\} \end{aligned}$$

and

$$\begin{aligned} & \int_0^1 x^{b-1}(1-x)^{a-2b+2} \\ & \times {}_7F_6 \left[ \begin{matrix} a, & \frac{1}{2}(a+4), & c, & d, & 2a-b-c-d+n+5, & 2-n, & 1 \\ \frac{1}{2}(a+2), & 3+a-c, & 3+a-d, & b+c+d-a-n-2, & a+n+1, & 3 \end{matrix} ; x \right] dx \\ & = \frac{(a-2)(b-a-2)_2(c-a-2)_2(d-a-2)_2(-a-n)_n(3+n+a-b-c-d)_2}{(a+2)(1-a)_2(1-b)_2(1-c)_2(1-d)_2(n-1)_2(b+c+d-2a-4-n)_2} \\ & \cdot \frac{\Gamma(b)\Gamma(a-2b+3)}{\Gamma(a-b+2)} \times \left\{ \frac{(a-1)_n(a-b-c+3)_n(a-b-d+3)_n(a-c-d+3)_n}{(a-b+1)_n(a-c+1)_n(a-d+1)_n(a-b-c-d+5)_n} \right. \\ & \quad \left. + \frac{na(b-2)(c-2)(d-2)(2a-b-c+n+3)}{(a-b+1)(a-c+1)(a-d+1)(b+c+d-a-n-4)(n+a-1)} - 1 \right\}. \end{aligned}$$

Similarly other results can be obtained.

### Concluding Remarks

In this paper, eleven Eulerian's type integrals involving generalized hypergeometric functions have been evaluated in terms of gamma function by employing very recently obtained summation theorems by Masjed-Jamei and Koepf. Several new, interesting and elementary integrals have also been given as special cases of our main findings.

### REFERENCES

- [1] G. E. ANDREWS, R. ASKEY and R. ROY, *Special Functions. In Encyclopedia of Mathematics and Its Applications, Volume 71*, Cambridge University Press, Cambridge, UK, (1999).
- [2] G. ARFKEN, *Mathematical Methods for Physicists*, Academic Press, New York, NY, USA, (1985).
- [3] W. N. BAILEY, *Generalized Hypergeometric Series*, Cambridge University Press, Cambridge, 1935; Reprinted by Stechert-Hafner, New York, (1964).

- [4] YU. A. BRYCHKOV, Evaluation of Some Classes of Definite Integrals, *Integral Transforms Spec. Fun.*, 13, pp. 163-167, (2012).
- [5] J. CHOI and A. K. RATHIE, Evaluation of Certain New Class of Definite Integrals, *Integral Transforms Spec. Fun.*, 22(4), pp. 282-294, (2015).
- [6] Y. S. KIM, M. A. RAKHA and A. K. RATHIE, Extensions of certain classical summation theorems for the series  ${}_2F_1$ ,  ${}_3F_2$  and  ${}_4F_3$  with applications in Ramanujan's summations, *Int. J. Math., Math. Sci.*, Article ID 309503, 26 pages, (2010).
- [7] W. KOEPF, *Hypergeometric Summation: An Algorithmic Approach to Summation and Special Function Identities*, 2nd ed., Springer, London, UK, (2014).
- [8] J. L. LAVOIE, F. GRONDIN and A. K. RATHIE, Generalizations of Watson's theorem on the sum of a  ${}_3F_2$ , *Indian J. Math.*, 34, pp. 23-32, (1992).
- [9] J. L. LAVOIE, F. GRONDIN, A. K. RATHIE and K. ARORA, Generalizations of Dixon's Theorem on the Sum of a  ${}_3F_2(1)$ , *Math. Comp.*, 62, pp. 267-276, (1994).
- [10] J. L. LAVOIE, F. GRONDIN and A. K. RATHIE, Generalizations of Whipple's theorem on the sum of a  ${}_3F_2$ , *J. Comput. Appl. Math.*, 72, pp. 293-300, (1996).
- [11] M. MASJED-JAMEI and W. KOEPF, Some Summation Theorems for Generalized Hypergeometric Functions, *Axioms*, 7, 38, 20 pages, (2018).
- [12] A. M. MATHAI and R. K. SAXENA, *Generalized Hypergeometric Functions with Applications in Statistics and Physical Sciences*, Lecture Note in Mathematics, Springer: Berlin/Heidelberg, Germany; New York, NY, USA, (1973); Volume 348.
- [13] A. F. NIKIFOROV and V. B. UVAROV, *Special Functions of Mathematical Physics. A Unified Introduction with Applications*, Birkhauser: Basel, Switzerland, (1988).
- [14] A. P. PRUDNIKOV, YU. A. BRYCHKOV and O. I. MARICHEV, *More Special Functions Integrals and Series*, vol. 3, Gordon and Breach Science Publishers, Amsterdam, The Netherlands, (1990).
- [15] E. D. RAINVILLE, *Special Functions*, The Macmillan Company, New York, 1960 ; Reprinted by Chelsea Publishing Company, Bronx, New York, (1971).
- [16] M. A. RAKHA and A. K. RATHIE, Generalizations of classical summation theorems for the series  ${}_2F_1$  and  ${}_3F_2$  with applications, *Integral Transforms Spec. Fun.*, 22(11), pp. 823-840, (2011).
- [17] L. J. SLATER, *Confluent Hypergeometric Functions*, Cambridge University Press, Cambridge, UK, (1960).
- [18] L. J. SLATER, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, UK, (1966).
- [19] H. M. SRIVASTAVA and J. CHOI, *Zeta and q-zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, (2012).